

PAJ282Hu01

Polyclonal Antibody to DNA Damage Inducible Transcript 3 (DDIT3)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µL

Cross Reactivity: Mouse; Rat

Applications: WB; IHC

### [ IMMUNOGEN ]

Immunogen: Recombinant DDIT3 (Met1~Ala169 (Accession # P35638)) expressed in E.coli

Accession No.: RPJ282Hu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

# Coud-Clone Corp.

expiration date under appropriate storage condition.

## [ IDENTIFICATION ]






Figure. Western Blot; Sample: Recombinant DDIT3, Human.



Vector Red staining on IHC-P;
Samples: Human Colon Tissue;
Primary Ab: 15µg/ml Rabbit AntiHuman DDIT3 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody



Vector Red staining on IHC-P;
Samples: Human Liver Tissue;
Primary Ab: 15µg/ml Rabbit AntiHuman DDIT3 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.