

| PA | N | 57  | 6H | h | n | 2 |
|----|---|-----|----|---|---|---|
| _  |   | .,, | v) |   | w | _ |

Polyclonal Antibody to Fibronectin Type III Domain Containing Protein 5 (FNDC5)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.44mg/ml

UOM: 1ml

Cross Reactivity: Mouse; Rat; Pig

Applications: WB; IHC; ICC; IP.

## [ IMMUNOGEN ]

Immunogen: Recombinant FNDC5 (Leu77~Met144 (Accession # Q8NAU1)) expressed in

E.coli

Accession No.: RPN576Hu02

#### [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

# Cloud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]





Figure. Western Blot; Sample: Recombinant FNDC5, Human.



DAB staining on IHC-P; Samples:
Human Cardiac Muscle Tissue;
Primary Ab: 20?g/ml Rabbit AntiHuman FNDC5 Antibody Second Ab:
2µg/mL HRP-Linked Caprine AntiRabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.