



A92472Hu03 Polyclonal Antibody to Fibulin 1 (FBLN1) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

| [PRODUCT INFORMATION] | Western Blot | | |
|--|--------------|---|--------|
| Immunogen: FBLN1 | kDa | | |
| Clonality: Polyclonal | 75 | | |
| Host: Rabbit | 50 | | |
| Species Reactivity: Human | 37 | | |
| Ig type: Rabbit IgG | 25 | | |
| Purification: Antibodies are purified by | 20 | - | ←24kDa |
| target protein affinity chromatography. | | | |
| Applications: WB, ICC, IHC-P, IHC-F | 15 | | |
| Form: Liquid | 10 | | |
| Size: 100µg | | | |

Sample: Recombinant human FBLN1

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant human FBLN1 (Asp399~IIe578) expressed in E.coli.

Molecular Weight: 21.4kDa

USCN accession No.: P92472Hu03

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-DV NECQRYPGRL CGHKCENTLG SYLCSCSVGF RLSVDGRSCE DINECSSSPC SQECANVYGS YQCYCRRGYQ LSDVDGVTCE DIDECALPTG GHICSYRCIN IPGSFQCSCP SSGYRLAPNG RNCQDIDECV TGIHNCSINE TCFNIQGGFR CLAFECPENY RRSAATLQQE KTDTVRCI

Unique product $\frac{S}{S}$ uperb quality Client favorite Nicest service (@ ISO9001:2008; @ ISO13485:2003; $\zeta \in$





[ANTIBODY SPEFICITY]

Anti FBLN1 is a rabbit polyclonal antibody raised against human FBLN1. This antibody has been selected for its ability to recognize human FBLN1 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant human FBLN1 (Asp399~IIe578) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Unique product Superb quality Client favorite Nicest service Q ISO9001:2008; Q ISO13485:2003; $\zeta \in$