

PAA321Hu01

Polyclonal Antibody to Galectin 1 (GAL1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

**UOM:** 100µl

Cross Reactivity: Mouse

**Applications:** WB,IHC

#### [ IMMUNOGEN ]

Immunogen: Recombinant GAL1 (Met1~Asp135) expressed in E.coli

Accession No.: RPA321Hu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

# Cloud-Clone Corp.

expiration date under appropriate storage condition.

### [ IDENTIFICATION ]







Figure. Western Blot; Sample: Recombinant GAL1, Human.



DAB staining on IHC-P;
Samples: Human Rectum Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody



DAB staining on IHC-P;
Samples: Human Renal cancer Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;


(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Rat Testis Tissue;
Primary Ab: 10µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Sample: Rat Colon Tissue;
Primary Ab: 10µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Rat Lung Tissue;
Primary Ab: 10µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

# Cloud-Clone Corp.




DAB staining on IHC-P;
Sample: Porcine Colon Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Porcine Uterus Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Porcine Small intestine Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Cardiac Muscle
Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Porcine Stomach Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman GAL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

kDa 1 2 3 4

116—
70—
44—
33—
27—
22—
18—
14—
10—

Western Blot; Sample: Lane1: K562 cell
lysate; Lane2: Mouse Heart lysate;
Lane3: Mouse Liver lysate; Lane4:
Mouse Kidney lysate
Primary Ab: 1µg/ml Rabbit Anti-Human
GAL1 Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

### [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was



used in clinical diagnostic or any other procedures.