



pA92053Mu01 Polyclonal Antibody to High Mobility Group AT Hook Protein 2 (HMGA2) Organism: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

7th Edition (Revised in May, 2013)

Western Blot

[PRODUCT INFORMATION]

Immunogen: HMGA2, Mouse

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100μg Sample: Recombinant HMGA2, Mouse



[IMMUNOGEN INFORMATION]

Immunogen: Recombinant HMGA2 (Met1~Asp108) expressed in *E.coli*.

USCN Accession No.: rP92053Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-MSARGEGAGQ PSTSAQGQPA
APVPQKRGRG RPRKQQQEPT CEPSPKRPRG RPKGSKNKSP SKAAQKKAET
IGEKRPRGRP RKWPQQVVQK KPAQETEETS SQESAEED





[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against HMGA2. It has been selected for its ability to recognize HMGA2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant HMGA2 (Met1~Asp108) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.