

PAA798Hu01

Polyclonal Antibody to Hypoxia Inducible Factor 1 Alpha (HIF1a)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 10µL

Cross Reactivity: Mouse; Porcine

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant HIF1a (Arg575~Asn826) expressed in E.coli

Accession No.: RPA798Hu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no


# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


### [ IDENTIFICATION ]




DAB staining on IHC-P; Sample: Mouse
Cardiac Muscle Tissue; Primary Ab:
20ug/ml Rabbit Anti-Human HIF1a
Antibody Second Ab: 2µg/mL HRPLinked Caprine Anti-Rabbit IgG
Polyclonal Antibody (Catalog:
SAA544Rb19)




DAB staining on IHC-P;
Samples: Human Cardiac Muscle
Tissue;
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Sample: Mouse Kidney Tissue;
Primary Ab: 20ug/ml Rabbit AntiHuman HIF1a Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Mouse Cardiac Muscle Tissue;
Primary Ab: 20ug/ml Rabbit AntiHuman HIF1a Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Sample: Porcine Colon Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal



DAB staining on IHC-P;
Sample: Porcine Small intestine Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal

# Coud-Clone Corp.



DAB staining on IHC-P;
Sample: Porcine Cardiac Muscle
Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked

(Catalog: SAA544Rb19)

Caprine Anti-Rabbit IgG Polyclonal

Antibody




Western Blot; Sample: Rat Serum
Primary Ab: 2µg/ml Rabbit Anti-Human
HIF1a Antibody

Second Ab: 0.2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)


Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Kidney Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)

Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Stomach Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman HIF1a Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)



## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.