

PAB837Po01

Polyclonal Antibody to Interleukin 2 Receptor Alpha (IL2Ra)
Organism Species: Sus scrofa; Porcine (Pig)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: IL2Ra, Pig Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



Sample: Recombinant IL2Ra, Pig

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant IL2Ra (Gly22~Gln245) expressed in E.coli.

Accession No.: RPB837Po01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- GACVQQPPS LRNATFKILG YKVGTTLNCD CQRGFRRDPS SGPYMICRGN SSHSFWENKC QCMPTSSPRI PVKQVTPRPE EQKERKTTET QGQMQPPNQA NLPGHCKEPP PWEHESLKRV YHFMEGQTVR YQCLPGFRDG SAQNNSAQSV CKKQEDQEVM RWTQPKLKCK SEKENGSEPE POMSTAAPPT TKTSI PTRTK GTTDSONI TE VPATMOPIIF TTQYQ



## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against IL2Ra. It has been selected for its ability to recognize IL2Ra in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant IL2Ra (Gly22~Gln245) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.