

PAB267Mu01

Polyclonal Antibody to Kininogen 1 (KNG1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.26mg/mL

**UOM:** 100µL

Cross Reactivity: N/A

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant KNG1 (Glu22~Arg379) expressed in E.coli

Accession No.: RPB267Mu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no



obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ IDENTIFICATION ]





DAB staining on IHC-P; Sample: Mouse
Kidney Tissue Primary Ab: 10µg/ml
Rabbit Anti-Mouse KNG1 Antibody
Control: Used PBS instead of primary
antibody Second Ab: 2?g/ml HRPLinked Caprine Anti-Rabbit IgG
Polyclonal Antibody (Catalog:

SAA544Rb19)

Western Blot; Samples: Lane1: Rat Serum; Lane2: Mouse Serum; Primary Ab: 0.05µg/ml Rabbit Anti-Mouse KNG1 Antibody Second Ab: 0.2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.