



A91490Hu01 Polyclonal Antibody To Peptidyl Prolyl Cis/Trans Isomerase NIMA Interacting Protein 4 (PIN4) Organism: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

[PRODUCT INFORMATION]

Immunogen: PIN4, Human

Clonality: Polyclonal


Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant PIN4, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant PIN4 (Ser23~Arg130) expressed in E.coli.

USCN Accession No.: P91490Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF- SADKKAQG PKGGGNAVKV RHILCEKHGK IMEAMEKLKS GMRFNEVAAQ YSEDKARQGG DLGWMTRGSM VGPFQEAAFA LPVSGMDKPV FTDPPVKTKF GYHIIMVEGR





[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against PIN4. It has been selected for its ability to recognize PIN4 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant PIN4 (Ser23~Arg130) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.