

PAA573Hu01

Polyclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)

Organism Species: *Homo sapiens* (Human)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Traits: Liquid

Concentration: 0.5mg/ml

UOM: 200µl

Cross Reactivity: Porcine

Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant PIIINP (Gln24~Pro153) expressed in *E.coli*

Accession No.: RPA573Hu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]







Figure. Western Blot; Sample: Recombinant PIIINP, Human.




DAB staining on IHC-P;
 Samples: Human Liver cancer Tissue;
 Primary Ab: 20µg/ml Rabbit Anti-Human PIIINP Antibody
 Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
 (Catalog: SAA544Rb19)




DAB staining on IHC-P;
 Sample: Human Liver Tissue
 Primary Ab: 20µg/ml Rabbit Anti-Human PIIINP Antibody
 Control: Used PBS instead of primary antibody
 Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
 (Catalog: SAA544Rb19)



DAB staining on IHC-P;
 Sample: Human Liver cancer Tissue
 Primary Ab: 20µg/ml Rabbit Anti-Human PIIINP Antibody
 Control: Used PBS instead of primary antibody
 Second Ab: 2µg/ml HRP-Linked




DAB staining on IHC-P;
 Sample: Human Placenta Tissue
 Primary Ab: 10µg/ml Rabbit Anti-Human PIIINP Antibody
 Control: Used PBS instead of primary antibody
 Second Ab: 2µg/ml HRP-Linked




DAB staining on IHC-P;
 Sample: Porcine Small intestine Tissue
 Primary Ab: 10µg/ml Rabbit Anti-Human PIIINP Antibody
 Control: Used PBS instead of primary antibody
 Second Ab: 2µg/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Sample: Porcine Cardiac Muscle
Tissue
Primary Ab: 10µg/ml Rabbit Anti-
Human PIIINP Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Testis Tissue
Primary Ab: 10µg/ml Rabbit Anti-
Human PIIINP Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Lane1: Hela cell
lysate; Lane2: HT1080 cell lysate
Primary Ab: 0.2?g/ml Rabbit Anti-
Human PIIINP Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

[**IMPORTANT NOTE**]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.