

PAA573Hu01 Polyclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP) Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [PRODUCT INFORMATION]

Immunogen: PIIINP-OVA

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 500µg/mL

#### **UOM:** 500µg

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Synthetic Peptide, PIIINP conjugated to OVA. Accession No.: CPA573Hu21 Sequence: The target peptide sequence is listed below. RDVWKPEPCQICVCD

#### [RELEVANCE]

PIIINP is the amino terminal peptide of type III procollagen, released from the precursor peptide during the synthesis and deposition of type III collagen. There is evidence that serum PIIINP measurement is an effective non-invasive test for the detection and monitoring of methotrexate-induced liver fibrosis and cirrhosis, and serial measurements may reduce the need for liver biopsy. PIIINP has a molecular weight of 42 000 and contains three distinct domains: a triple-helical domain (Col 3) in the middle of the molecule, the Col 1-domain at the amino terminal and Col 2-domain at the carboxyterminal end of the propeptide.

Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003 11271 Richmond Avenue Suite H104, Houston, TX 77062, USA | Tol free: 001-888-960-7402 | Fax: 001-832-538-0068 | Http://www.doud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F. Wuhan, Hubel 430056, PRC | Tol free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Http://www.uscnk.com | E-mail: mail@uscnk.com

## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against PIIINP conjugated to OVA. It has been selected for its ability to recognize PIIINP in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

# [ QUALITY CONTROL ]

Content: The quality control contains synthetic peptide PIIINP.Usage: Please dissolve the synthetic peptide PIIINP in water before use.Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.