

CSI087Gu01

Primary Guinea Pig Adult Dermal Fibroblasts (ADF)

Organism Species: Cavia (Guinea pig)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Revised in May, 2025)

### [ DESCRIPTION ]

**Cell Type:** Fibroblast **Synonyms:** ADF

Species: Cavia (Guinea pig)

Tissue Source: Adult Dermal tissue

**Disease:** Normal **Size:** >5×10<sup>5</sup>cell/vial

### [ PROPERTIES ]

Cell activity: >85% (Viability by Trypan Blue Exclusion).

Formulation: Frozen 1 mL or T25 flask.

Biosafety: Negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.

Applications: For research use only. It is not approved for Feline or animal use, or for application in

clinical diagnostic procedures. **Growth Properties:** Adherent

# [CONTENTS]

Form & Buffer: Supplied as solution form in frozen stock solution, containing 90% FBS+10% DMSO.

# [USAGE]

Upon receiving the cells in a T-25 flask at room temperature, immediately transfer the cells to 37°C, 5% CO<sub>2</sub> incubator; the cells in vials, directly and immediately transfer the cells from dry ice to liquid nitrogen.

#### **Culture conditions:**

DMEM+10%FBS+ 1%Fibroblast growth supplement +1%Penicillin-Streptomycin Solution

Temperature: 37°C

Condition: 95% air, 5% carbon dioxide

#### Cell recovery:

After receiving the cells, shake at 37°C in a water bath until completely dissolved, transfer to a 15 ml centrifuge tube, add 3-5 times complete culture solution, 1000 rpm for 5 min, discard the supernatant, and place in a T25 flask for culture.

#### Cell passage:

1. Cell passage when cell growth at 85-95%.



- 2. Discard the medium and wash with PBS 1-2 times.
- 3. Add 1 ml of Trypsin at 37°C, observe the cell under the microscope. If the cells are retracted and rounded, pat the culture flask to let the cells fall off. Stop digestion by adding 2 ml of complete medium containing 10% serum. Make it a single cell suspension.
- 4. Add the fresh medium to resuspend the cells. Unless otherwise stated, the recommended ratio of primary cells is 1/2.

# [Shipping]

Dry ice.

## [STORAGE]

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

## [IMPORTANT NOTE]

The cell is for research use only, and we will not be responsible for any issue if the cell was used in clinical diagnostic or any other procedures.

# [Figure]




Figure 1 Figure 2

Figure 1 Morphology of Guinea Pig Adult Dermal Fibroblasts (Optical microscope, x100) Figure 2 Morphology of Guinea Pig Adult Dermal Fibroblasts (Optical microscope, x200)