

RPF182Hu01 200µg Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 12 (HSD17b12) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Asn312

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 120µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.3

Predicted Molecular Mass: 38.0kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

Con <u>Cloud-Clone Corp</u>.

MESALPAAGF LYWVGAGTVA YLALRISYSL FTALRVWGVG NEAGVGPGLG EWAVVTGSTD GIGKSYAEEL AKHGMKVVLI SRSKDKLDQV SSEIKEKFKV ETRTIAVDFA SEDIYDKIKT GLAGLEIGIL VNNVGMSYEY PEYFLDVPDL DNVIKKMINI NILSVCKMTQ LVLPGMVERS KGAILNISSG SGMLPVPLLT IYSATKTFVD FFSQCLHEEY RSKGVFVQSV LPYFVATKLA KIRKPTLDKP SPETFVKSAI KTVGLQSRTN GYLIHALMGS IISNLPSWIY LKIVMNMNKS TRAHYLKKTK KN

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.