

RPD088Mu01 100µg

Recombinant Acid Phosphatase 6, Lysophosphatidic (ACP6)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Met1~Glu418

Tags: N-terminal His Tag

Subcellular Location: Mitochondrion

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.6

Predicted Molecular Mass: 51.3kDa

Accurate Molecular Mass: 51kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MRVWVPVGV L TSLAYCFHQ R VALAEQRAP NGQRPVDRNL LELKMVQVVF RHGARSPLKP LPLEEQVEWN  
PKLLEIPPQT RFDYTVT NLA GGPKPHSHYD TEYRKTTLRG GVLAGQLTKV GMQQMFALGE KLRKKNYVEDI  
PFLSPVYNPQ EVFIRSTNMF RNLESTRCLL AGLFQHQKGS AVIHTDEASS EVLYPNYQSC WVLKEKTRGR  
KKA AISQPGI SEDLEKVK TG VGINNGDDVD FVLLDNVAA EQVHSLNCP ALERFAQLIE QRAVDMALYV  
VEQEDRESIQ MAVGPF LHL EGNLLKT VDP TTAPSKTRKM YLYATHDVTL LPMLLALGIF DQKWPPFAVD  
LTMELYQHQE SKEW FVQLFY NGKEQVPRGC PDKLCPLDKF LNTMSVYSVS PEKYRTLCSK TRIVELGE
```

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.