

RPA824Hu01 10μg Recombinant B-Lymphocyte Activation Antigen B7-2 (LAB7-2) Organism Species: *Homo sapiens (Human) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Coud-Clone Corp.

## [PROPERTIES]

Source: Prokaryotic expression Host: *E.coli* 

Residues: Gln46~Cys282

Tags: N-terminal His Tag

Subcellular Location: Membrane

**Purity:** > 95%

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.2

Predicted Molecular Mass: 30.9kDa

Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.

#### [<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


#### [SEQUENCE]

# Cloud-Clone Corp.


QNQSL SELVVFWQDQ ENLVLNEVYL GKEKFDSVHS KYMGRTSFDS DSWTLRLHNL QIKDKGLYQC IIHHKKPTGM IRIHQMNSEL SVLANFSQPE IVPISNITEN VYINLTCSSI HGYPEPKKMS VLLRTKNSTI EYDGVMQKSQ DNVTELYDVS ISLSVSFPDV TSNMTIFCIL ETDKTRLLSS PFSIELEDPQ PPPDHIPWIT AVLPTVIICV MVFCLILWKW KKKKRPRNSY KC

# [IDENTIFICATION]





### Figure. Gene Sequencing (Extract)



### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.