

RPC410Hu01 100µg Recombinant Cathepsin E (CTSE) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Ser20~Pro396

Tags: N-terminal His and GST Tag

Subcellular Location: Endosome

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.5

Predicted Molecular Mass: 70.7kDa

Accurate Molecular Mass: 71kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[<u>SEQUENCE</u>]

	S	LHRVPLRRHP	SLKKKLRARS	QLSEFWKSHN
LDMIQFTESC S	MDQSAKEPL	INYLDMEYFG	TISIGSPPQN	FTVIFDTGSS
NLWVPSVYCT S	PACKTHSRF	QPSQSSTYSQ	PGQSFSIQYG	TGSLSGIIGA
DQVSVEGLTV V	/GQQFGESVT	EPGQTFVDAE	FDGILGLGYP	SLAVGGVTPV
FDNMMAQNLV D	LPMFSVYMS	SNPEGGAGSE	LIFGGYDHSH	FSGSLNWVPV
TKQAYWQIAL D	NIQVGGTVM	FCSEGCQAIV	DTGTSLITGP	SDKIKQLQNA
IGAAPVDGEY A	VECANLNVM	PDVTFTINGV	PYTLSPTAYT	LLDFVDGMQF
CSSGFQGLDI H	PPAGPLWIL	GDVFIRQFYS	VFDRGNNRVG	LAPAVP

[IDENTIFICATION]




Figure. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.