

**RPB463Ra01 200µg**

**Recombinant Chitinase-3-like Protein 1 (CHI3L1)**

**Organism Species: *Rattus norvegicus* (Rat)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

12th Edition (Revised in Aug, 2016)

## [ **PROPERTIES** ]

**Source:** Prokaryotic expression

**Host:** *E.coli*

**Residues:** Tyr117~Lys364

**Tags:** N-terminal His Tag

**Subcellular Location:** Nucleus, Cytoplasm, Extracellular matrix

**Purity:** > 95%

**Traits:** Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

**Original Concentration:** 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 8.7

**Predicted Molecular Mass:** 31.2kDa

**Accurate Molecular Mass:** 29kDa as determined by SDS-PAGE reducing conditions.

## [ **USAGE** ]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## [ **STORAGE AND STABILITY** ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ **SEQUENCE** ]

YGFD  
GLDLAWLYPG PKDKQHFTTL IKELKAEFTK EVQPGTEKLL LSAAVSAGKV  
TLDSGYDVAQ IAQHLDIFINL MTYDFHGTWR HTTGHHSPLF RGQQDTGPDR  
FSNVDYGVGY MLRLGAPTNK LVMGIPTFGK SFTLASSENQ VGAPITGSGL  
PGRYTKEKGT LAYYEICDFL RGAEVHRILG QQVPFATKGN QWVGYYDDPES  
VKNKVLYLKN KQLAGAMVWA VDLDDFRGSF CGHNVHFPLT NAIK

**[ IDENTIFICATION ]**




Figure. SDS-PAGE

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.