# Cloud-Clone Corp.

**RPA988Hu01 10µg**

**Recombinant Cytochrome P450 2E1 (CYP2E1)**

**Organism Species: *Homo sapiens (Human)***

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

## [ PROPERTIES ]

**Source:** Prokaryotic expression

**Host:** *E.coli*

**Residues:** Pro36~Val271

**Tags:** N-terminal His and GST Tag

**Subcellular Location:** Membrane, Endoplasmic reticulum lumen

**Purity:** > 95%

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

**Original Concentration:** 200μg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 9.3

**Predicted Molecular Mass:** 57.6kDa

**Accurate Molecular Mass:** 58kDa as determined by SDS-PAGE reducing conditions.

## [ USAGE ]

Reconstitute in ddH$_2$O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8⁰C for one month.

Aliquot and store at -80⁰C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCE ]

```
                                    PFPLP IIGNLFQLEL
KNIPKSFTRL AQRFGPVFTL YVGSQRMVVM HGYKAVKEAL LDYKDEFSGR
GDLPAFHAHR DRGIIFNNGP TWKDIRRFSL TTLRNYGMGK QGNESRIQRE
AHFLLEALRK TQGQPFDPTF LIGCAPCNVI ADILFRKHFD YNDEKFLRLM
YLFNENFHLL STPWLQLYNN FPSFLHYLPG SHRKVIKNVA EVKEYVSERV
KEHHQSLDPN CPRDLTDCLL V
```

## [ IDENTIFICATION ]



**Figure . Gene Sequencing (extract)**



Figure. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.