

RPF192Hu01 1 Recombinant Fucosyltransferase 2 (FUT2) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Gln29~His343

Tags: N-terminal His Tag

Subcellular Location: Membrane, Golgi apparatus

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.7

Predicted Molecular Mass: 39.4kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCE</u>]

Cond-Clone Corp.

QQ RLAKIQAMWE LPVQIPVLAS TSKALGPSQL RGMWTINAIG RLGNQMGEYA TLYALAKMNG RPAFIPAQMH STLAPIFRIT LPVLHSATAS RIPWQNYHLN DWMEEEYRHI PGEYVRFTGY PCSWTFYHHL RQEILQEFTL HDHVREEAQK FLRGLQVNGS RPGTFVGVHV RRGDYVHVMP KVWKGVVADR RYLQQALDWF RARYSSLIFV VTSNGMAWCR ENIDTSHGDV VFAGDGIEGS PAKDFALLTQ CNHTIMTIGT FGIWAAYLTG GDTIYLANYT LPDSPFLKIF KPEAAFLPEW TGIAADLSPL LKH

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.