1304 Langham Creek Dr. Suite 226, Houston, TX 77064, USA | 001-888-960-7402 | www.choud-clone.us | mail/d cloud-clone.as Expurt Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-clour.com | mail/u-cloud-clour

RPC110Hu01 100µg **Recombinant Growth Differentiation Factor 5 (GDF5) Organism Species: Homo sapiens (Human)** Instruction manual

### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES


[PROPERTIES] Residues: Ala382~Arg501 Tags: N-terminal His-Tag Accession: P43026 Host: E. coli Subcellular Location: Secreted. **Purity:** >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in 10mM PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative. 15% SDS-PAGE Predicted isoelectric point: 6.5 Predicted Molecular Mass: 15.1kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

# [USAGE]

Reconstitute in sterile ddH<sub>2</sub>O.

10th Edition (Revised in Jan, 2014)

kDa



Coud-Clone Corp.

# Cloud-Clone Corp.

### Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [ <u>SEQUENCES</u> ]

The sequence of the target protein is listed below.

APLATRQGK RPSKNLKARC SRKALHVNFK DMGWDDWIIA PLEYEAFHCE GLCEFPLRSH LEPTNHAVIQ TLMNSMDPES TPPTCCVPTR LSPISILFIDSANNVVYKQY EDMVVESCGC R

## [REFERENCES]

- 1. Polinkovsky A., et al. (1997) Nat Genet 17 (1): 18-9.
- 2. Thomas JT., et al. (1997) Nat Genet 17 (1): 58-64.
- 3. O'Keeffe G., et al. (2004) J Neurocytol 33 (5): 479-88.
- 4. Buxton P., et al. (2001) J Bone Joint Surg Am 83-A Suppl 1 (Pt 1): S23-30.
- 5. Francis-West P., et al. (1999) Development 126 (6): 1305–15.