RPA074Rb61 100uq Recombinant Immunoglobulin G1 (IgG1) Organism Species: Oryctolagus cuniculus (Rabbit) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Glu1~Lys180 Accession: P01870 Host: 293F cell **Purity: >95%** Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in 20mM Tris,

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 8.0

Predicted Molecular Mass: 25.0kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note: The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.


5. Polymerization of the target protein: Dimerization, multimerization etc.

1304 Langham Creek Dr, Suite 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-clone.us | mail@cloud-clone.us Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-800-880-0687 | www.cloud-clone.com | mail@cloud-clone.com

kDa 70 44 33 26 22 18 14

15% SDS-PAGE

10



Coud-Clone Corp.

Coud-Clone Corp.

[<u>USAGE</u>]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

EDDPEVQFTW YINNEQVRTA RPPLREQQFN STIRVVSTLP IAHQDWLRGK EFKCKVHNKA LPAPIEKTIS KARGQPLEPK VYTMGPPREE LSSRSVSLTC MINGFYPSDI SVEWEKNGKA EDNYKTTPAV LDSDGSYFLY SKLSVPTSEW QRGDVFTCSV MHEALHNHYT QKSISRSPGK