

RPB629Hu01 50µg

Recombinant Insulin Like Growth Factor 2 Receptor (IGF2R)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Asn2167~Ala2442

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.6

Predicted Molecular Mass: 31.6kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
NGDN YLYEIQLSSI TSSRNPACSG ANICQVKPND
QHFSRKVGTS DKTKYYLQDG DLDVVFASSS KCGKDKTKSV SSTIFFHCDP
LVEDGIPEFS HETADCQYLF SWYTSAVCPL GVGFDSENPG DDGQMHKGLS
ERSQAVGAVL SLLLVALTCC LLALLLYKKE RRETVISKLT TCCRRSSNVS
YKYSKVNKEE ETDENETEWL MEEIQLPPPR QGKEGQENGH ITTKSVKALS
SLHGDDQDSE DEVLTIPEVK VHSGRGAGAE SSHPVRNAQS NA
```

[IDENTIFICATION]

\$MITOMISSSOW_DICTIONICION_THE TO DICKMS_TO SMC033351S_TEGSSOWAINS CASSISWASCOWNSHOWS CONTROSCOS WAS REACTED AND CONTROSCOS CONTROSCO





Figure . Gene Sequencing (extract)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.