

RPA895Hu02 50 μ g
Recombinant Insulin Receptor (ISR)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Arg1027~Met1364

Tags: N-terminal His-Tag

Tissue Specificity: Heart, Kidney, Spleen.

Subcellular Location: Membrane; Single-pass type I membrane protein.

Purity: >92%

Traits: Freeze-dried powder

Buffer formulation: 10mM PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 39.6kDa

Accurate Molecular Mass: 39kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```

RELG QGSFGMVYEG NARDIIKGEA
ETRVAVKTVN ESASLRERIE FLNEASVMKG FTCHHVVRLL GVSKGQPTL
VVMELMAHGD LKSYLRSLRP EAENNPGRPP PTLQEMIQMA AEIADGMAYL
NAKKFVHRDL AARNCMVAHD FTVKIGDFGM TRDIYETDYY RGGKGLLPV
RWMAPESLKD GVFTTSSDMW SFGVVLWEIT SLAEQPYQGL SNEQVLKFVM
DGGYLDQPDN CPERVTDLMR MCWQFNPKMR PTFLEIVNLL KDDLHPSFPE
VSFFHSEENK APESEELEME FEDMENVPLD RSSHCQREEA GGRDGGSSLG
FKRSYEEHIP YTHM
    
```

[**IDENTIFICATION**]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE