

RPC044Hu01 200µg

Recombinant Liver X Receptor Alpha (LXRa)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



# [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Asn95~Lys434

**Tags:** N-terminal His Tag

**Subcellular Location:** Nucleus

**Purity:** > 90%

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 100µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 41.0kDa

**Accurate Molecular Mass:** 41kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ SEQUENCE ]

```
NELCSV
CGDKASGFHY NVLSCEGCKG FFRRSVIKGA HYICHSGGHC PMDTYMRRKC
QECRLRKCRQ AGMREECVLS EEQIRLKKLK RQEEEQAHAT SLPPRASSPP
QILPQLSPEQ LGMIEKLVAA QQQCNRRSFS DRLRVTPWPM APDPHSREAR
QQRFAHFTEL AIVSVQEIVD FAKQLPGFLQ LSREDQIALL KTSAIEVMLL
ETSRRYNPGS ESITFLKDFS YNREDFAKAG LQVEFINPIF EFSRAMNELQ
LNDAEFALLI AISIFSADRP NVQDQLQVER LQHTYVEALH AYVSIHHPHD
RLMFPRMLMK LVSLRTLSSV HSEQVFALRL QDKK
```

## [ IDENTIFICATION ]





Figure . Gene Sequencing (extract)



## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.