

RPG790Hu02 50µg

Recombinant Mdm2 p53 Binding Protein Homolog (MDM2)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Met1~Pro491

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.6

Predicted Molecular Mass: 58.9kDa

Accurate Molecular Mass: 80kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MCNTNMSVPT DGAVTTSQIP ASEQETLVRP KPLLLKLLKS VGAQKDTYTM
KEVLFYLQY IMTKRLYDEK QQHIVYCSND LLGDLFGVPS FSVKEHRKIY
TMIYRNLVVV NQQESSDSGT SVSENRCHLE GGSDQKDLVQ ELQEEKPSSS
HLVSRPSTSS RRAISETEE NSDELSGERQ RKRHKSDSIS LSFDESLALC
VIREICCERS SSSESTGTPS NPDLDAVSE HSGDWLDQDS VSDQFSVEFE
VESLDESDYS LSEEGQELSD EDDEVYQVTV YQAGESDTDS FEEDPEISLA
DYWKCTSCNE MNPPLPSHCN RCWALRENWL PEDKGKDKGE ISEKAKLENS
TQAEEGFDVP DCKKTIVNDS RESCVEENDD KITQASQSQE SEDYSQPSTS
SSIIYSSQED VKEFEREETQ DKEESVESSL PLNAIEPCVI CQGRPKNCCI
VHGKTGHLMA CFTCAKLLKK RNKPCPVCRQ PIQMIVLTYF P
```

[IDENTIFICATION]

G A A T T C A T G T G C A A T A C C A A C A T G T C T G T A C C T A C T G A T G G T G C T G T A A C C A C C T C A C A G A T T C C A G C T T C G G A A C A A G A G A C C T G G T T A G A C C A A G C C A T T G C T T T T G A A G T T A T T A A A G T C T G T T G G T G C A C A A A A G A C A




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.