

RPB202Hu01 10µg

Recombinant Neurokinin B (NKB)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residuess: Cys23~Glu121

Tags: N-terminal His Tag

Tissue Specificity: Secreted

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.1

Predicted Molecular Mass: 14.8kDa

Accurate Molecular Mass: 17kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

CKEPQEEV VPGGGRSKRD PDLYQLLQRL
 FKSHSSLEGL LKALSQASTD PKESTSPEKR DMHDFVGLM GKRSVQPDSP
 TDVNQENVPS FGILKYPPRA E

[IDENTIFICATION]

JGTAAAGGACCAAGGAGGAGGTGCTCTGCGGCGCCGCAAGAGGGATCCAGATCTTAAGCTCTCCAGAGCTCTCAAAAGCCACTCATCTCGAGGATCTCAAGCCCTGAGCCAGCCAGCATCTAGGATCAACATCTCCGGAAGGTGACATGCATGACTCTTTGTGGCTTATGGCAA
 C K E P Q E E V V P G G G R S K R D P D L Y Q L L Q R L F K S H S S L E G L L K A L S Q A S T D P K E S T S P E K R D M H D F F V G L M G K




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE