

**RPB388Ca01 10 $\mu$ g**

**Recombinant Neutrophil Gelatinase Associated Lipocalin (NGAL)**

**Organism Species: *Canis familiaris*; Canine (Dog)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## **[ PROPERTIES ]**

**Residues:** Ser23~Glu198

**Tags:** N-terminal His-Tag

**Accession:** E2RSM4

**Host:** *E. coli*

**Purity:** >90%

**Endotoxin Level:** <1.0EU per 1 $\mu$ g

(determined by the LAL method).

**Formulation:** Supplied as lyophilized form in 10mM

PBS, pH7.4, containing 1mM DTT, 5% trehalose,


0.01% sarcosyl and preservative.

**Predicted isoelectric point:** 7.1

**Predicted Molecular Mass:** 21.7kDa

**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



## **[ USAGE ]**

Reconstitute in sterile ddH<sub>2</sub>O.

## **[ STORAGE AND STABILITY ]**

**Storage: Avoid repeated freeze/thaw cycles.**

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The sequence of the target protein is listed below.

STPSLIPA PPPLKVPLQP DFQHDQFQGK WYVIGIAGNI LKKEGHGQLK MYTTTYELKD  
DQSYNVTSTL LRNERCDYWN RDFVPSFQPG QFSLGDIQLY PGVQSYLVQV VATNYNQYAL  
VYFRKVYKSQ EYFKITLYGR TKELPLELKK EFIRFAKSIG LTEDHIIFPV PIDQCIDE

## **[ REFERENCES ]**

1. Lindblad-Toh K., *et al.* (2005) Nature 438:803-819.
2. Yang J., *et al.* (2002) Mol. Cell 10 (5): 1045–56.
3. Flo TH., *et al.* (2004) Nature 432 (7019): 917–21.
4. Schmidt-Ott KM., *et al.* (2007) J. Am. Soc. Nephrol. 18 (2): 407–13.