

RPC828Hu01 10 μ g
Recombinant Tripeptidyl Peptidase I (TPP1)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Met1~Pro320

Tags: N-terminal His-Tag

Tissue Specificity: Heart, Placenta.

Subcellular Location: Lysosome. Melanosome.

Purity: >92%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.7

Predicted Molecular Mass: 38.2kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MRLFGGNFAH QASVARVVGQ QGRGRAGIEA SLDVQYLMSA GANISTWVYS
SPGRHEGQEP FLQWLMLLSN ESALPHVHTV SYGDDEDSLS SAYIQRVNTE
LMKAAARGLT LLFASGDSGA GCWSVSGRHQ FRPTFPASSP YVTTVGGTSF
QEPFLITNEI VDYISGGGFS NVFPRPSYQE EAVTKFLSSS PHLPPSSYFN
ASGRAYPDVA ALSDGYWVVS NRVIPWVSG TSASTPVFGG ILSLINEHRI
LSGRPPLGFL NPRLYQQHGA GLFDVTRGCH ESCLDEEVEG QGFCSGPGWD
PVTGWGTPNF PALLKTLNLP

[IDENTIFICATION]




Figure 1. SDS-PAGE