

RPL817Hu01 10µg

Recombinant Wingless Type MMTV Integration Site Family, Member 4 (WNT4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Leu26~Arg351

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.8

Predicted Molecular Mass: 40.0kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[SEQUENCE]

		LYLAK	LSSVGSISEE	ETCEKLKGLI
QRQVQMCKRN	LEVMDSVRRG	AQLAIEECQY	QFRNRRWNCS	TLDSLPVFGK
VVTQGTREAA	FVYAISSAGV	AFAVTRACSS	GELEKCGCDR	TVHGVSPQGF
QWSGCSDNIA	YGVAFSQSFV	DVRERSKGAS	SSRALMNLHN	NEAGRKAILT
HMRVECKCHG	VSGSCEVKTC	WRAVPPFRQV	GHALKEKFDG	ATEVEPRRVG
SSRALVPRNA	QFKPHTDEDL	VYLEPSPDFC	EQDMRSGVLG	TRGRTCNKTS
KAIDGCELLC	CGRGFHTAQV	ELAERCSCKF	HWCCFVKCRQ	CORLVELHTC
R				

[IDENTIFICATION]




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.