

RPA128Mu01 100μg
Tissue Inhibitors Of Metalloproteinase 2 (TIMP2)
Organism Species: Mus musculus (Mouse)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Cys27~Pro220 (Accession # P25785),
with N-terminal His-Tag.

Host: *E. coli*

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg
(determined by the LAL method).


Formulation: Supplied as lyophilized form in PBS,
pH7.4, containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 6.5

Predicted Molecular Mass: 23.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.
MGHHHHHHSG SEF-CSCS PVHPQQAFCN ADVVIRAKAV SEKEVDSGND IYGNPIKRIQ
YEIKQIKMFK GPDKDIEFIY TAPSSAVCGV SLDVGGKKEY LIAGKAEGDG KMHITLCDFI
VPWDTLSITQ KKSLNHRYQM GCECKITRCP MIPCYISSPD ECLWMDWVTE KSINGHQAKF
FACIKRSDGS CAWYRGAAPP KQEFLDIEDP

[REFERENCES]

1. Ma S., *et al.* (2013) PLoS Biol. 11:e1001469-e1001469.
2. Mi M., *et al.* (2012) Biochem. Biophys. Res. Commun. 423:366-372.
3. Lee Y., *et al.* (2011) Dev. Cell 21:746-757.
4. Kandam V., *et al.* (2011) Circulation 124:2094-2105.