

RPA139Mu01 10µg

Recombinant Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## [PROPERTIES]

Residues: Met40~Asn291

**Tags:** Two N-terminal Tags, His-tag and S-tag

Accession: P50592

Host: E. coli

Subcellular Location: Membrane; Single-pass

type II membrane protein.

**Purity: >90%** 

**Endotoxin Level:** <1.0EU per 1μg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris,

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% sarcosyl, 5% trehalose, and preservative.




Predicted Molecular Mass: 35.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

# [USAGE]

Reconstitute in ddH<sub>2</sub>O.





### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [SEQUENCES]

The sequence of the target protein is listed below.

M YFTNEMKQLQ DNYSKIGLAC FSKTDEDFWD STDGEILNRP CLQVKRQLYQ LIEEVTLRTF QDTISTVPEK QLSTPPLPRG GRPQKVAAHI TGITRRSNSA LIPISKDGKT LGQKIESWES SRKGHSFLNH VLFRNGELVI EQEGLYYIYS QTYFRFQEAE DASKMVSKDK VRTKQLVQYI YKYTSYPDPI VLMKSARNSC WSRDAEYGLY SIYQGGLFEL KKNDRIFVSV TNEHLMDLDQ EASFFGAFLI N

#### [REFERENCES]

- 1. Wiley S.R., et al. (1995) Immunity 3:673-682.
- 2. Cretney E., et al. (2002) J. Immunol. 168:1356-1361.
- 3. Takeda K., et al. (2002) J. Exp. Med. 195:161-169.
- 4. Strebel A., et al. (2002) Int. J. Cancer 100:627-634.