

**RPE929Hu01 50µg**

**Recombinant Vav 3 Oncogene (VAV3)**

**Organism Species: Homo sapiens (Human)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## **[ PROPERTIES ]**

**Residues:** His183~Glu380

**Tags:** N-terminal His-Tag

**Accession:** Q9UKW4

**Host:** *E. coli*

**Purity:** >95%

**Endotoxin Level:** <1.0EU per 1µg

(determined by the LAL method).


**Formulation:** Supplied as lyophilized form in 10mM PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative.

**Predicted isoelectric point:** 6.4

**Predicted Molecular Mass:** 24.7kDa

**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



15% SDS-PAGE

## **[ USAGE ]**

Reconstitute in sterile ddH<sub>2</sub>O.

## **[ STORAGE AND STABILITY ]**

**Storage: Avoid repeated freeze/thaw cycles.**

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The sequence of the target protein is listed below.

HQPKCPEN DIRSCCLAEI KQTEEKYTET LESIEKYFMA PLKRFLTAAE FDSVFINIPE  
LVKLHRNLMQ EIHSIVNKN DQNLVQVFIN YKERLVIYQG YCSGVESAIS SLDYISKTK  
DVKLKLEECs KRANNGKFTL RDLLVVPMPQRLVKYHLLLQE LVKHTTDPTE KANLKLALDA  
MKDLAQYVNE VKRDNETLRE

## **[ REFERENCES ]**

1. Movilla N., Bustelo X.R. (1999) Mol. Cell. Biol. 19:7870-7885.
2. Zeng L., *et al.* (2000) Mol. Cell. Biol. 20:9212-9224.
3. Yabana N., Shibuya M. (2002) Oncogene 21:7720-7729.
4. Hunter S.G., *et al.* (2006) Mol. Cell. Biol. 26:4830-4842.